



Welcome to All Creatures Great and Small!

Your puppy's vaccine program will depend on the age and previous vaccine history. This is some information on K-9 vaccines and testing.

Yes	No	Cost	<u>K-9 Distemper vaccine</u>
<input type="checkbox"/>	<input type="checkbox"/>	\$35	<p>K-9 Distemper, also known as Distemper/Parvo or DA2P-P, is a combination vaccine that starts at 6 weeks of age, and boosters are given 3-4 weeks apart until 16-18 weeks of age to protect against the following diseases:</p> <ul style="list-style-type: none"> • Distemper: a viral disease that is often fatal in puppies. It is characterized by respiratory distress, coughing, and seizures; it is highly contagious. • Hepatitis (Adenovirus): a viral disease spread through urine. It causes respiratory and liver problems. • Parainfluenza: a respiratory infection that is often involved with kennel cough. • Parvovirus: a severe intestinal viral disease characterized by vomiting and bloody diarrhea. It is often fatal and highly contagious.
Yes	No	Cost	<u>Rabies vaccine</u>
<input type="checkbox"/>	<input type="checkbox"/>	\$35	<p>The rabies vaccine is given when the puppy is 12 weeks or older. Rabies is a serious and fatal disease that attacks the brain and the central nervous system. It is most dangerous and is a zoonotic disease. The vaccine is made from a killed rabies virus. The first vaccine lasts for 1 year. Then, it is given every three years.</p>
Yes	No	Cost	<u>Bordetella vaccine</u>
<input type="checkbox"/>	<input type="checkbox"/>	\$35	<p>Bordetella is also known as kennel cough or tracheobronchitis. The bordetella vaccine is started at 10 weeks of age, and boosters are given 3-4 weeks apart and then annually after. It is highly contagious and often difficult to treat.</p>
Yes	No	Cost	<u>Lyme vaccine</u>
<input type="checkbox"/>	<input type="checkbox"/>	\$37	<p>The Lyme vaccine starts at 10 weeks or older, and boosters are given 3-4 weeks apart and then annually thereafter. The Lyme vaccine protects against Lyme disease, which is caused by a bacteria called <i>Borrelia burgdorferi</i>, which is passed to dogs through a bite from a deer tick. Lyme disease can cause lameness, kidney failure, and heart disease. This is a dangerous zoonotic disease.</p>
Yes	No	Cost	<u>K-9 Influenza vaccine</u>
<input type="checkbox"/>	<input type="checkbox"/>	\$40	<p>K-9 influenza vaccine, or CIV, can start at 8 weeks of age or older, and boosters are given 2-3 weeks apart and then annually thereafter. K-9 influenza causes a respiratory infection in dogs.</p>
Yes	No	Cost	<u>Leptospirosis vaccine</u>
<input type="checkbox"/>	<input type="checkbox"/>	\$40	<p>The leptospirosis vaccine starts at 12 weeks or older, and boosters are given 3-4 weeks apart and then annually thereafter. Leptospirosis is a disease caused by infection by <i>Leptospira</i> bacteria. <i>Leptospira</i> bacteria can be found worldwide in soil and water. Leptospirosis can cause flu-like symptoms, liver disease, and kidney disease. It is a zoonotic disease.</p>

Yes No Cost
 \$62

Heartworm, Lyme, Anaplasma, and Ehrlichia Testing

Heartworm & tick-borne diseases testing is recommended starting at 6 months of age. Heartworm disease can be fatal and is transmitted by infected mosquitoes. Once the dog has been infected, heartworm larvae make their way to the dog's heart and pulmonary arteries.

Yes No Cost
 Costs will vary

Heartworm Prevention

Heartworm prevention can begin at 8 weeks of age and comes as a monthly pill.

Yes No Cost
 Costs will vary

Vaccine Titrers

Vaccine titers are blood tests performed to check your dog's level of immune defenses instead of vaccinating. This is usually an option after the series of puppy vaccines are finished.

Yes No Cost
 \$65

Ova, Parasites, & Giardia Testing

Having your dog's stool checked is very important, and it is recommended that there be at least two negative samples for puppies. Most puppies have worms. The most common ones are hookworms, tapeworms, roundworms, whipworms, and coccidian. All worms attack the intestines.

Yes No Cost
 Costs will vary

Neuters & Spays

Neutering and spaying your puppy is important and recommended at 6 months of age or older. Your pet will generally have fewer health problems and a better temperament in the future.

Yes No Cost
 \$85
(includes registration)

Microchipping

A microchip is inserted under your dog's skin. This is not painful and can give you peace of mind if your pet is lost and put in a shelter. We recommend microchipping at time of spay or neuter.

Yes No Cost
 \$25

Ear Mites

Ear mites are parasites located in the ears. Ear mites can cause ear and skin diseases, which can spread to other pets. Having an ear swab done can determine whether your dog is infected with ear mites.

Yes No Cost
 Costs will vary

Flea & Tick Prevention

Flea & tick prevention is highly recommended. There are topical flea & tick preventatives that come as a liquid that is applied to your dog's skin. There are also oral flea & tick preventatives. Flea & tick preventatives can be started at 8 weeks of age or older.

Signature

Date